

Ch #12 136

An Essay  
on Passed March 24  
1825

Acute Peritonitis

by  
Gustavus B. Campbell

of Virginia

of which  
greater  
that he  
on it  
as an in  
to mount  
has obse  
througho  
of the pe  
the conse  
is the  
incident  
as, & ag  
hardly  
deviation  
two he  
The lo  
demon  
the tena  
degrees 7

Peritonitis is an inflammatory affection of the Peritoneum; that serous membrane which affords a lining to the parietes of the abdomen, as well as an exterior coat & a means of attachment to its viscera. It is, also, attended by general febrile symptoms; & it has, therefore, all the essential features, which entitle a disease to a place in the Order Phlegmasia & Clap Pyrexia of Cullen.

Peritonitis is divided into the Acute & the Chronic. It is the Acute species, that constitutes the subject of this Thesis; & my remarks will be restricted to that form of the disease, to which both sexes are rendered liable by the operation of common causes.

History of Acute Peritonitis. An attack of Acute Peritonitis usually commences with rigors. When reaction takes place, there is fever, marked by thirst; a dry state of the tongue & fauces; a pulse, accelerated, corded, & contracted. The patient soon feels a considerable degree of pain & heat in the abdomen;

insupportable  
 heats from  
 the sun, the  
 quickness  
 that cannot  
 back, no  
 dry & hot,  
 vomiting,  
 the face  
 the tongue  
 the appetite  
 lips, & gums  
 the stools  
 constant  
 punctions.  
 Unusually  
 low. The  
 inclination  
 to  
 tons are  
 of pain,  
 almost ce  
 so rapid



& when pressure is made on that part, manifests a greater or lesser degree of tenderness, which may either be limited to a part of the abdomen, or spread over it universally. Though the tongue & fauces are in general dry, as already stated, it is proper to mention, that in some rare instances, they have been observed to retain their natural moisture throughout the course of the disease. The state of the pulse also is subject to great variety at the onset of the disease: while in most cases, it possesses the character of inflammatory action, to a very evident degree; in others, it is scarcely disturbed; & again, in others it is so low, as to be hardly perceptible. The cases which present these deviations of the pulse, will be particularly noticed hereafter.

The local uneasiness increases rapidly, & the abdomen swells. Within twelve or twenty-four hours, the tenderness of the abdomen is raised to such a degree, that the weight of the bed-clothes is then

*[Faint, illegible handwriting, likely bleed-through from the reverse side of the page.]*

offer ge  
held cas  
counten  
helped or  
acted or  
upon the  
- A  
On dispo  
of the Re  
entire.  
thus are  
this is  
of infla  
canal, on  
has inve  
also. Th  
are mu  
have. O  
the by  
jukes in  
regulate  
are bud  
tens, cl

intolerable. The abdomen is now tense. The pulse beats from a hundred & twenty, to a hundred & thirty strokes in a minute. The respiration much quickened. The strength, much depressed. The patient cannot bear the least motion; but lies on his back, with the knees drawn up. The skin is dry & hot, though in instances of much nausea & vomiting, or of great pain, it is occasionally damp. The face is flushed, & expresses anxiety. The eyes are red. The tongue is white, as well as dry. The thirst is great. The appetite is diminished. And the patient is restless, & gets but little sleep.

The Stomach, in most cases, is much affected; & a constant sickness, with a vomiting of bile, is experienced. — The intestines, sometimes costive; at others, preternaturally loose; are at others still perfectly regular. The bladder, also, is affected, & there is a constant inclination to pass urine, attended however with little effect.

As the disease advances, the symptoms are all aggravated. At length, a total cessation of pain suddenly takes place; an omen, which indicates, almost certainly, a fatal issue. The pulse sinks, but moves so rapidly as not to be counted. No dark matter, like

the Periton  
Prognosis  
abdomen  
other of op  
in the con  
some pulle  
cating, eas  
ally, the p  
is affected  
disease.

Diaon  
by the abdo  
the lower pa  
sustaining  
It is distin  
by the plac  
of special  
trinitis, ab  
drawn up,  
Carles  
to Perito  
which reg  
les, chemo

coffee grounds, is ejected from the stomach by singultus. Cold clammy sweats break out. The extremities are cold. The countenance is collapsed & haggard. The urine & feces are passed involuntarily. And difficult respiration is succeeded by the fatal event, which usually takes place before the eighth day.

#### - Appearances on Dissections -

On dissection, the marks of inflammation appear, in every part of the Peritonæum; with the exception, as has been stated, of the anterior. The Peritonæum is thicker, & less transparent. Its arteries are injected with red blood, in most cases, but occasionally, there is found no vestige of the kind. Commonly, there are no traces of inflammation in the abdominal muscles: in the intestinal canal, on the contrary, it is sometimes seen, that the inflammation has involved, not only the muscular, but the more interior coats also. The sides of this canal, & the duplications of Peritonæum, are much thickened by extravasation into their cellular membrane. The viscera of the abdomen are agglutinated to one another by coagulating lymph. A brownish fluid, like serum, is effused into the Peritonæal cavity; & has mixed with it, shreds of coagulating lymph, & sometimes pus. Marks of gangrene however are but seldom visible. These appearances, independently of symptoms, clearly shew that the disease is an inflammation of

State of  
Alabama  
alone or c  
sequence of

The paper  
much dispe  
the large on  
the price of  
thought of a  
appears to  
dispute, re  
tion.

The plan  
more stron  
it is not  
the more  
ful sanc  
on of the c  
bity of c

the Peritoneum.

**Prognosis.** — This is unfavorable, when the pain in the abdomen suddenly ceases; the pulse sinks; & there is evidence either of effusion, or of gangrene. The following appearances on the contrary, are to be regarded as favorable. The pulse becomes fuller, softer & less frequent; the skin moist & cool; the respiration easier; the pain & tension of the abdomen, diminished gradually; the patient, able to maintain a sitting posture; & the bowels, affected with a gentle diarrhoea towards the close of the disease.

**Diagnosis.** Peritonitis is distinguished from Enteritis, by the absence of that tenesmus, which exists in Enteritis, when the lower part of the intestines are inflamed; & by the local pain sustaining no relief from fecal evacuations.

It is distinguished from Colic; by the absence of tormina; by the pain being augmented, rather than abated, by pressure; & especially, by the difference of posture; the patient, in Peritonitis, always choosing to lie on his back, with his knees drawn up; & in Colic, on his abdomen. —

**Causæ.** Among the causes which give a predisposition to Peritonitis, are, habits of intemperance; occupations which require the body to be habitually bent, as that of tailors, shoemakers & wash women; & a habitual exposure to cold.

*[Faint, mirrored handwriting, likely bleed-through from the reverse side of the page.]*

Unsuspected  
picked  
its employ  
it produces  
that. The  
stage is a  
by which  
The justice  
truly by 1.  
the pulse  
dibility, p  
avoids. If  
judging of  
upward to  
portion A  
at their en  
by the de  
toms.  
If the pa  
of blood.  
any sense  
movements  
peated a



Acute Peritonitis is excited by acts of violence, done to the abdomen; but oftener, by Cold applied for some time, either alone or combined with moisture. Thus it may be the consequence of sleeping either in damp sheets or on the ground.

### Treatment

The proper treatment of Peritonitis has been a subject for much dispute. The Medical School of England, recommends the large employment of Opium. Dr. Brunus of Dublin employs the Spirit of Turpentine, in Purpurul Peritonitis, at the very height of inflammation. While others, selecting a plan, which appears to be obviously, deduced from the pathology of the disease, rely for its cure chiefly on abundant depletion.

The plan of treatment now to be detailed, is to myself more strongly recommended than any other: not only is it in my own opinion well suited to the case; but in its more important points at least, it derives a powerful sanction from the experience of the present Professor of the Institutes & Practice of Physic in the University of Pennsylvania.



*Venesection.* In acute Peritonitis, the lancet is to be considered as the patient's sheet anchor; & in the extent of its employment, it is to be regulated by the effects which it produces on the disease, & on the constitution of the patient. The appearance of debility existing in the early stage is deceptive, & depends on the violence of the attack, by which the powers of the system are oppressed.

The justice of this remark will be illustrated satisfactorily, by the effect of a liberal detraction of blood; as the pulse will then be found to rise, & the apparent debility, previously existing, will be proportionally removed. It is the advice of Dr. Nathaniel Chapman, that in judging of the propriety of Venesection, we should pay no regard to the pulse, which is always depressed in proportion to the violence of the disease. I should be governed then entirely, by the existence of the stage of excitement, & by the degree of severity in the local pain & other symptoms.

If the patient be an adult, take at first  $\mathfrak{z}\text{xx}$  or  $\mathfrak{z}\text{xxx}$  of blood. If this detraction should <sup>produce</sup> on the symptoms no very sensible effect; or if that effect were evidently an improvement, the operation may be advantageously repeated within a few hours.

had a ten-  
towards  
have comba-  
not tend  
property of  
in of dispo-  
sibility of or  
conclude a  
which it  
Bostwick's  
is it to  
good to the  
of De. (Mon-  
abdomen,  
by proceed  
the guar-  
ally, by s-  
on three p-  
collisions  
open; but  
they may  
Wash-  
nothing;

Topical Blood-letting—Topical bloodletting is next to be resorted to; & is performed by means of leeches or of cups applied to the abdomen. As in local inflammations generally, a liberal abstraction of blood in this way will be very useful. It has been conceived that this measure ought to be deferred, until the pain of the abdomen has been somewhat diminished; or at least, until the constitutional effects of the local derangement have been partially removed, by Venesections. But I should not suppose that the trivial irritation of the measure, would produce any serious effect on the general excitement; & I should be unwilling, for the sake of preventing a slight effect of the kind, to forego the probable advantage to be obtained from an early use of the remedy.

Purgatives. Immediately after Venesection, I would administer a dose of Calomel, which the torpid state of the bowels generally requires to be large. The dose may be gr. XX or even half a Drachm. This is to be followed by an infusion of Senna with Sattu, or Epsom Salts alone, & administered every hour, until several copious evacuations have been obtained. If the stomach be so irritable as to reject the Calomel, a large Cathartic enema may be administered. Evacuating the lower part of the intestines

the disease  
a bag -  
they show  
not as 7  
disappear  
is the end  
of St. Ch.  
It does a  
it oaks  
Children  
this purp  
would a  
is com  
the mean  
all the sh  
dory, it  
dancy a  
I would  
there or p  
Naticate  
but are to  
at the cas  
by wear

has a tendency to favour the retention of purgatives administered by the mouth. The Calomel may also have combined with it a little Opium, which here does not tend to constipate.

I know that the propriety of active purging in acute Peritonitis, is a matter of dispute. It appears to me however, that, independently of other reasons, the acknowledged utility of the measure in all other cases of Phlegmasia; & the service which it has done in the analogous case of Pleurisy Peritonitis, are fully sufficient to warrant us in extending it to the case under consideration.

With regard to the use of purging, I would adopt here, the rule of Dr. Armstrong: "So long as there are pain & tension of the abdomen, a quick pulse, & offensive stools, I would steadily proceed." While the bowels were not under the operation of a purgative, I would keep them soluble, by small doses of saline mixture given every two or three hours.

Emollient Enemata will assist in keeping the bowels open; but are useful chiefly as internal fomentations they may consist of warm water, or flaxseed tea.

Warm fomentations are in their immediate effect soothing; & ultimately make a great impression on

anger -

to check or

the Epineu-

reidarium,

Dist. -

stage of Accu-

istic. It

early war

Thud s

let an reg

its stage of

efforts affo

low, or sh

disease. -

operations of

lax to dig

is the s

in this case

But is

the person



the disease. Among the best are, a bread & milk poultice; a bag of muslin; or flannels wrung out of hot water. They should be so large as to cover the abdomen, & as hot as the patient can bear them.

Diaphoretics. next in importance to bloodletting is the inducing of perspiration. In the strong language of Dr. Chapman, this measure often acts like a charm. It does good here by the centrifugal determination which it gives to the blood; thus relieving the vessels of the Peritoneum of a portion of their burden. For this purpose, I would employ the Vapour Bath; & would assist its action by some one of the Diaphoretics combined with Opium. Previously to depletion, by the means already recited, Opium would aggravate all the symptoms; but after this has been sufficiently done, it will do good, as well by its diaphoretic tendency as by relieving pain & morbid vigilance. I would employ the Dover's Powder  $\text{Dose } \frac{ij}{\text{gr.}}$  every three or four hours.

Vesication. Blisters, well timed are of much advantage; but are too irritating to be employed in the commencement of the case. They should be preceded by copious depletion & by warm fomentations. When applied, they should be

ed however  
of the  
Should acc  
directly pro  
we should  
vision, for  
one of the a  
specific blue  
measures, we  
complete re  
sultion. —  
tendency of  
you has app  
and shows ho  
likely the ste  
able. — I can  
house of this c  
very eminent  
will present as  
of historical

large.

To relieve irritation of the stomach, & to check bilious vomiting, mucilaginous drinks are advised, & the Effervescent draught, with fifteen or twenty drops of laudanum, if necessary, may be exhibited.

Diet. — It is unnecessary to say that the diet, in the first stage of Acute Peritonitis, ought to be strictly antiphlogistic. It should consist of such articles as toast-water, barley-water, rice-water &c.

Thus have I detailed all the important measures, that are required for the treatment of Acute Peritonitis in its stage of Excitement. It is in this stage only that our efforts afford any prospect of success; & in this stage, therefore, we should bring all our resources to bear upon the disease. —

When the pulse sinks, & the other symptoms of Collapse supervene; depletion should give place to diffusible stimulants. The best of these probably is the spirits of Turpentine. Dr. Chapman says that in this case, he has repeatedly seen it of decided utility. But it is necessary to give it in the dose of a <sup>tea</sup> ~~table~~ spoonful, every two or three hours. It is to be a pist.



ed however, with Vol. alkali & wine; & with Sinapisms  
& Blisters to the extremities.

Should Acute Peritonitis be marked at its onset, by a pulse  
scarcely perceptible, & by other signs of extreme prostration;  
we should consider it as a case, not of exhaustion, but of op-  
pression; for it owes its peculiar character entirely to the vio-  
lence of the disease. Here, we are first to make use of small,  
repeated bleedings, the Warm bath, & Diaphoretics, When by these  
measures, we have roused the energies of the system, & established  
a complete reaction, we may freely resort to more copious de-  
pletion. — Peritonitis sometimes begins with slight

tenderness of the abdomen, the pulse scarcely affected, very little  
fever, & no appearance whatever calculated to alarm. But the se-  
quel shews how insidious is this aspect. For suddenly, the pulse  
sinks, the strength fails, & the case proceeds rapidly to a fatal  
close. — A case of this kind occurred last summer in the Almshouse  
of this city. It was mistaken by the prescribing physician,  
a very eminent practitioner, for simple Constipation. I was my-  
self present at the post mortem examination; when the evidence  
of Peritoneal inflammation was found to be very satisfactory,

the as few  
with that p  
which he has  
exclusively  
one of the  
house has  
particular  
on the sta  
but there  
with an app  
and, in  
important, and  
inflammation  
in house a  
up to the  
and sometimes  
tion of a lo  
the self- and  
the abdom  
the cases of